

Search for linear measures to estimate the anatomical composition of the carcass

J. NAVEAU

*I.T.P., Centre expérimental de Sélection porcine,
Maxent, 35380 Plélan-le-Grand (France)*

The purpose of this study was to verify in French conditions, the conclusions of Professor de Boer's report concerning estimation of carcass composition by simple linear measures.

Measurements were made according to de Boer's method on 323 pig carcasses in the Stations of Mauron and Le Rheu.

The results obtained are in agreement with those of Zeist. Prediction equations of muscle and fat weight of the carcass were established. The accuracy of these equations was $R^2 = 0.82$ for muscle weight and $R^2 = 0.85$ for fat weight.

Estimation of the anatomical composition of pig carcasses and of different cuts

I. — Establishment of fatness indexes

II. — Classification according to indexes

B. DESMOULIN ⁽¹⁾, P. GRANDSART ⁽²⁾ et J. P. VILA ⁽³⁾

⁽¹⁾ *Station de Recherches sur l'Élevage des Porcs
I.N.R.A.-C.N.R.Z., 78350 Jouy-en-Josas (France)*

⁽²⁾ *I.T.P.-M.N.E., 149, rue de Bercy, 75579 Paris Cedex 12*

⁽³⁾ *Laboratoire de Biométrie
I.N.R.A.-C.N.R.Z., 78350 Jouy-en-Josas (France)*

The present study deals with the estimation of carcass fatness in 126 pigs of various genotypes, sexes and nutritional statuses. Eight sites of backfat thickness (easily approachable variables) were used to establish fatness indexes supplying information about the accurate references of the overall fat content (uneasily approachable variables).

1) In practice, optimum prediction of fatness by means of a simplified procedure involve use of either two lateral measures taken at a distance of 4, 5 and 8 cm from the vertebral column (index $1.65 P_1 + P_3$) or two measures on the split at the level of loin and neck (index $1.08 ER^3 + 1.06 EC$). The specificity of the indexes was due to their high prediction value of the weight and distribution of fat.

2) Segmentation of the variation of each index into 5 classes allowed to establish a classification according to the index in close relation with the fatness references of each group; the latter were characterized by the uneasily approachable measures (backfat weight, flare-fat, specific gravity of belly). The linear segmentation of the indexes might be further changed in order to obtain the best within-group homogeneity. The limits between classes established according to indexes represented a better sorting of fatness classes than that resulting from use of the loin/back maxima (CEE index).

3) For 2 classes of carcass weights (70-79 kg and 80-89 kg) the variation of measures and of fatness indexes was studied in relation with the limits of muscle content classes established by the CEE grade. The results obtained were the following :

a) Correction of backfat thickness in relation to $a + 10$ kg rise in carcass net weight varied according to the sites of measurements. It appeared to be irregular at the level of the neck and