

## The effect of age on fibre digestion in the rabbit

I. G. PARTRIDGE

*The Rowett Research Institute, Bucksburn, Aberdeen AB2 9SB  
(Great Britain)*

The ability of the commercial meat rabbit to digest dietary fibre was examined at three different ages (6 weeks, 14 weeks and 22 weeks). Dried grassmeal provided the principal fibre source in two experimental diets fed *ad libitum*. The ADF content of the diets was 130 g ADF/kg DM (diet A) and 226 g ADF/kg (diet B).

There was no apparent improvement in the DM digestibility of either diet as the animal aged. Overall the DM digestibility coefficients for diet A and diet B were 71 p. 100 and 54 p. 100 respectively. Similarly there was no improvement in ADF digestibility with age, coefficients of ADF digestibility being on average 13 p. 100 (diet A) and 12 p. 100 (diet B). Animals fed continuously on the high fibre diet did not consistently digest more fibre than those animals changed from the low to high fibre diet for the duration of each balance period, i.e. there was no evidence of animals adapting their fibre digesting capabilities. The low fibre digestibilities reported could be associated with the degree of lignification of these grassmeal-based diets (approx. 22 p. 100 of the ADF fraction).

---

## Effect of different crude fibre levels on nutrient digestibility of diets for growing rabbits

R. DESSIMONI CARREGAL and C. F. ZINSLY

*Faculdade de Ciências Agrárias e Veterinárias « Campus » de Jaboticabal  
UNESP — Dept. de Produção Animal. Rod. Carlos Tonanni s/n. Jaboticabal-SP  
(Brazil)*

The work was made at the Rabbit culture Section of Faculdade de Medicina Veterinária e Agronomia de Jaboticabal, with the aim to observe the effect of different crude fibre levels on the nutrient digestibility of growing rabbits as well as to estimate the digestion capacity of the fibrous fraction of the diet.

The trials were made with 15 New Zealand white rabbits using diets containing 7, 10 and 13 p. 100 crude fibre and the experiment was achieved with 5 replications entirely at random.

The following conclusions were drawn: 1 — the digestibility coefficient of organic matter, protein and non nitrogenous extract reduced with increasing levels of crude fibre in the diet. 2 — Rise in the crude fibre levels of the diet increased its digestibility coefficients. This increase was probably due to the analytical method used.

---

## Energy feeding of rabbits with low cost and high fibre materials

J. M. CID DIAZ

*Antonio Lopez 51, Madrid, 29 (Spain)*

Eight New-Zealand doe rabbits (8 month-old) and young rabbits of 2nd parturition were fed diets at two energy levels using low cost feeds without lucerne (2 900 kcal digestible energy (DE) per kg feed and 2 625 kcal DE/kg for doe-rabbits and 2 835 to 3 150 kcal/kg feed for their