

## Research on the use of dehydrated whole maize plant in the feed of growing rabbits

Maria-Teresa AUXILIA, G. MASOERO

*Istituto Sperimentale per la Zootecnia  
Sezione Operativa Periferica. V. Pianezza 115, Torino 10151 (Italy)*

Pelleted feeds containing 0 p. 100 or 20 p. 100 or 40 p. 100 dried whole-crop maize at the wax stage were given to growing rabbits.

A rearing trial was carried out in 52 rabbits weaned at 28 days of age and slaughtered at 84. The average daily gain was quite similar in the three groups, while the voluntary intake was increased and the feed efficiency was reduced with dried maize at 40 p. 100.

The digestibility coefficients of nutrients and fibrous components of polysaccharides in the three diets during three periods between 41 and the 70<sup>th</sup> days of age were tested.

The digestibility coefficients increased during the trial, but significantly until 60 days of age.

## Utilization of dehydrated whole maize plant for rabbits

Madeleine SEROUX <sup>(1)</sup>, Y. FRANCK <sup>(2)</sup>, P. MERCIER <sup>(3)</sup>

*(1) ITCF, Station Expérimentale, Boigneville, 91720 Maise (France).*

*(2) ITAVI, Service Technique, 28, rue du Rocher, 75008 Paris (France).*

*(3) UFAC, Service Cunicole, 95450 Vigny (France).*

Four trials were made to study dehydrated whole maize plant as crude fibre source in the diets of fattening rabbits.

Mature maize was incorporated into the diet at three levels: 30, 50 and 70 p. 100. Furthermore (except in one trial) two stages of harvest were compared, hard dough stage versus mature at the level of 50 p. 100 of the diet.

Crude protein and crude fibre contents of the feeds were 15 — 16 p. 100 and 14 — 15 p. 100. Neither the level of maize in the diet nor its stage of harvest affected the average daily gain.

A high proportion of maize at the latest harvest slightly reduced the feed consumption and accordingly the efficiency was improved.

Carcass yield was not modified.

## Utilization of dehydrated beet pulp for growing rabbits

Y. FRANCK <sup>(1)</sup>, Madeleine SEROUX <sup>(2)</sup>

*(1) ITAVI, 28, rue du Rocher, 75008 Paris (France).*

*(2) ITCF, Boigneville, 91700 Maise (France).*

Two trials were made to study the use of dehydrated sugar beet pulp in fattening rabbits (incorporation levels : 0 — 10 — 20 — 30 p. 100).

In the experimental conditions, and particularly with formulas including 16 p. 100 of Weende cellulose, no effect of high pulp levels on mortality was noticed.