

For the bacon pigs subjected to restricted feeding with diets containing 3 300 kcal D.E. supplying 2.50 to 2.60 g lysine for 1 000 kcal D.E. and offered in the trough as a moistened meal, the presence of 5 *points lupin* did not alter the performance as compared to the control diet only composed of maize and soybean meal. At the level of 10 *points*, a decrease in growth rate as well as a 4 p. 100 increase in the feed conversion ratio were observed over the whole fattening period. These discrepancies only occurred during the finishing period. At the level of 15 *points* the performance were highly reduced (by 10 p. 100) and the animals exhibited flatulence troubles accompanied by a reduced appetite. Use of flax seeds as curative or preventive treatment undoubtedly reduced the magnitude of these phenomena, which are probably due to the bad digestion of alpha-galactosides whose proportion is rather high in this type of lupin (8 p. 100 stachyose).

A reduction in the carcass yield was noticed in pigs fed lupin containing diets but it has to be pointed out that 10 and 15 points lupin led to an improvement of the body composition.

Nutritional value of whole extruded soybean seeds in piglet feeding

A. AUMAITRE, D. BOURDON

*I.N.R.A., Station de Recherches sur l'Élevage des Porcs,
Centre de Rennes-Saint-Gilles, F 35590 L'Hermitage*

Soybean oil-meal intended for monogastric animals has to be subjected to heat treatments so as to improve its nutritional value. Possible use of whole soybean seeds gives rise to a double problem involving the presence of antinutritional factors and a high lipid level. Moreover, only little accurate information is available on the possible use of such a feed in early weaning diets for piglets.

This experiment allowed to compare a classical diet composed of maize-soybean meal (70 and 22 p. 100) with a almost isoenergetic diet composed of barley, a moderately energy rich cereal associated with whole extruded soybean seed (67 and 25 p. 100). Piglet performance measured between 5 and 10 weeks of age showed that whole extruded seed which contained less than 0.8 ITU of antitrypsin factor per mg was as valuable a protein source as soybean meal. Both diets were equally consumed (866 versus 876 g/day) and the daily mean gain was similar for both treatments (444 versus 447 g/day). Feed conversion ratios were identical (1.95 versus 1.94 kg of diet per kg weight gain). A supplementary isoenergetic diet based on barley-soybean-meal and tallow led to equivalent mean performance.

However, the adaptation of the animals varied according to diet composition. Performances during the different periods between 8.4 and 24 kg were discussed. Conclusions were drawn as regards the efficiency of the extrusion in suppressing antinutritional factors and the economic conditions of the use of whole soybean seeds in weaning diets for piglets.

Choice of the protein sequence during post-weaning (*ad libitum*) and fattening periods (restricted)

F. GROSJEAN *, J. CASTAING **, O. LAVOREL *, M. LEUILLET *

* *I.T.C.F., 8, avenue du Président-Wilson, F 75116 Paris*

** *A.G.P.M., 122, boulevard Tourasse, F 64000 Pau*

Two experiments were carried out with 256 *Large White* pigs weighing from 9 to 103 kg.

In the first experiment we studied the subsequent effect of the post-weaning protein level (21 or 18 p. 100 crude protein) on bacon pigs receiving a suitable level of protein (18 p. 100). The following sequences were compared : 21-18 and 18-18.