

to karyotype determination (either on themselves or on their progeny), nearly 40 p. 100 were revealed carriers of a chromosomal abnormality. Translocations are by far the most frequent chromosomal abnormalities in the pig. A total of 22 different translocations have been so far identified in this species and among them, 6 were described in France. The decrease in size of the litters issued from breeding animals (males or females) carrying a translocation results from the production of gametes with an unbalanced karyotype (either gain or loss of chromosomal material) : these gametes are normally fertile but give non-viable embryos which disappear at the time of implantation. About one half of surviving foetuses are carriers of the translocation.

Among the six new translocations identified in France (some of them have been repeatedly found), three lead to a marked decrease in prolificacy : 49, 45 and 41 p. 100 for the 4-, 14+, 3+, 7- and 4-, 15+ translocations, respectively. The effect on litter size is of smaller magnitude for the others : 31, 28 and 26 p. 100 for the 16+, 17-, 5-, 14+ and 1-, 15+, respectively. Different proposals aiming at increasing precision and quickness of detection are discussed. The economic consequences of a translocation carried by a breeding boar vary according to the type of farm but are anyway of great importance. Finally, chromosomal abnormalities may be a valuable tool in several fields of research, such as genetic mapping (e.g. localization of the SLA complex on chromosome n° 7) and chromosome sorting by means of flux cytometry (creation of DNA banks for studies of molecular biology).

### **Evolution of litter size with the age of the dam at farrowing : Updating adjustment coefficients in the French pig breeds**

R. GUEBLEZ <sup>(1)</sup>, C. LEGAULT <sup>(2)</sup>, J. DAGORN <sup>(1)</sup>, Geneviève LE HENAFF <sup>(2)</sup>

*(1) Institut Technique du Porc, Région Ouest, B.P. 3, 35650 Le Rheu*

*(2) Institut National de la Recherche Agronomique, Station de Génétique quantitative et appliquée, 78350 Jouy-en-Josas*

The evolution of litter size with the age of the dam at farrowing was studied from data collected through the French National Litter Recording Programme in 1983-1984 for 5 breeds of sows : Large-White (112,886 litters), French Landrace (28,536 litters), Pietrain (1,189 litters), Belgian Landrace (4,660 litters) and sows used in the commercial units, mostly crossed Large-White × French Landrace sows (over one million litters). Polynomial (3rd degree) regression coefficients on the age of the dam at farrowing were calculated for various dependent variables : litter size at birth (total born or born alive) and at weaning, stillborn and mortality rate. These results were then compared to those from a similar study carried out 12 years before. In the Large-White and French Landrace, litter size increases with the age of the dam up to a plateau, but the curve is plateauing around 150 days earlier than 12 years ago, i.e. at around 850 days for litter size at birth and 650 days at weaning ; moreover this plateau is lower by around 0.75 total born or born alive, or 0.25 weaned piglets. This so-called decrease in litter size potential is to be connected with changes in management practices, particularly with a decrease of the age at weaning (from 42 down to 28 days over this period). We also observed that litter size generally tended to be higher in the Pietrain than in the Belgian Landrace. These results led to improved adjustment coefficients of litter size for the age of the dam. Various possibilities of using this correction were discussed.