

Intestinal fatty acid digestion and energy utilization in lambs infused with different plant oils into the abomasum

T Obitsu, H Ieki, K Taniguchi

Faculty of Applied Biological Science, Hiroshima University, Higashihiroshima-shi 724, Japan

Protected fat escaping ruminal degradation is regarded as a useful additive for the increase of energy supply. However, fat sources entering the small intestine may affect intestinal fatty acid digestion and nutrient utilization in host animal. The objective of this study was to investigate the effects of different plant oils infused into the abomasum on intestinal fatty acid digestion and energy utilization in lambs.

Three male lambs (initial body weight : 26.6 ± 2.1 kg) fitted with abomasal and ileal cannulae were used in a 3×3 Latin square design. Lambs were fed a basal diet (40 % hay and 60 % concentrate) at 1.6 times maintenance energy level. Emulsified coconut oil (CO), palm oil (PO) or rice bran oil (RO) with Cr-EDTA was continuously infused into the abomasum for 17 days of each period. The daily amount of fat infusion was 5 % (w/w) of the basal diet ($2.5 \text{ g/BW}^{0.75}$ day). As a control, the lambs were fed the basal diet with abomasal infusion of soy lecithin (an emulsifying reagent) for 17 days before the Latin square. Ileal digesta, feces, urine and venous blood were collected during the last 6 days of each period. Ileal digesta flow was calculated by the indicator method. Fatty acid contents in

ileal digesta and feces were measured by GLC after direct methylation (Sukhija and Palmquist, 1988, J Agric Food Chem, 36, 1202-1206). Heat production over 24 hours was measured by indirect calorimetry.

Gross energy intake (GE) tended to be lower for CO and PO than for control and RO due to the decreases in dry matter intakes. Energy digestibility (DE/GE) tended to be higher for CO than for PO. Heat production (HP) was not affected by fat infusion and fat sources. Energy retention to metabolizable energy (ER/ME) was higher for RO than for other three treatments. Total fatty acids (TFA) entering the ileum were variable, but fecal TFA excretion was lower for CO than for PO. Intestinal digestibility of infused TFA was 97.4, 86.8 and 93.7 % for CO, PO and RO, respectively. Plasma concentrations of triglycerides and non-esterified fatty acids tended to increase with fat infusion. Plasma total cholesterol was higher for CO and PO than for control.

These results indicated that energy of RO infused into the abomasum may be used more efficiently than that CO and PO in growing lambs.

Item	Control	CO	PO	RO	SEM
GE ($\text{kJ/kg}^{0.75}$)					
Basal diet	1121	977	969	1051	26
Basal diet + oil	1121	1092	1091	1175	26
DE/GE (%)	78.5	81.6	78.0	79.3	0.8
ME ($\text{kJ/kg}^{0.75}$)	745 ^{ab}	764 ^{ab}	736 ^a	804 ^b	6
HP ($\text{kJ/kg}^{0.75}$)	534	544	516	526	6
ER ($\text{kJ/kg}^{0.75}$)	212 ^a	220 ^a	220 ^a	278 ^b	3
ER/ME (%)	28.4 ^a	28.8 ^{ab}	29.7 ^{ab}	34.5 ^b	0.5
Total fatty acid flow					
Ileum ($\text{mg/kg}^{0.75}$)	277	130	718	392	85
Feces ($\text{mg/kg}^{0.75}$)	200 ^a	243 ^a	509 ^b	341 ^{ab}	24

a, b : $P < 0.05$.