

Erratum

Anim. Res. 54 (2005) 241–257

Effects of hay quality on intake, growth path, body composition and muscle characteristics of Salers heifers

Thierry HOCH^{a*}, Catherine JURIE^a, Philippe PRADEL^b,
Isabelle CASSAR-MALEK^a, Roland JAILLER^a, Brigitte PICARD^a,
Jacques AGABRIEL^a

^a Unité de Recherche sur les Herbivores, INRA, Theix, 63122 Saint-Genès-Champanelle, France

^b Domaine de Marcenat, INRA, 15190 Marcenat, France

(Received 14 June 2004; accepted 1 February 2005)

Due to misprints, Figures 1 and 2 have been badly printed. We reproduce below the correct figures.




Figure 1. Evolution of Full Body Weight (in kg) of continuously (thick line) or discontinuously (thin line) growing heifers, with standard error. For legibility, a two week step is used for data plotting.

* Corresponding author: hoch@vet-nantes.fr

Present address: UMR ENV Nantes INRA Gestion de la Santé Animale, Atlanpole-Chantrerie, BP 40706, 44307 Nantes Cedex 03, France.




Figure 2. Evolution of daily dry matter intake (in kg.d⁻¹) of continuously (thick line) or discontinuously (thin line) growing heifers, with standard error. For legibility, a two week step is used for data plotting.